

Magnetic resonance imaging of the heart: more than the morphological analysis

Viktor Peršić*

Thalassoterapija Opatija, Opatija, Croatia

Magnetic resonance imaging (MRI) of the heart is one of the key non-invasive techniques in modern cardiology. In the last period it developed from the useful research method into a clinically evidence-based, safe and comprehensive diagnostic test. The development of the technology has resulted in its wider application in various fields of cardiology, in evaluation of regional and global systolic function of the heart, perfusion and tissue characterization of the heart muscle, evaluation of pericardial disease, heart tumor and follow-up of patients with congenital heart disease and diseases of the aorta. In ischemic heart disease there are great possibilities for identifying segmental contractility failures af-

ter administration of dobutamine, or applying vasodilating tests such as adenosine, with a satisfactory safety profile and a good degree of diagnostic accuracy. The principle of gadolinium-based imaging is based on a lower speed of wash out of gadolinium from myocardium replaced by fibrosis or scar. On these grounds, a delayed imaging 5 to 20 minutes after injection of contrast agent will clearly show a fibrosis or a scar and thus also give an answer to the question as to the benefits of further revascularization procedures. The main advantages of cardiac MRI versus other non-invasive imaging methods are high spatial resolution, excellent reproducibility (suitable for monitoring patients), non-ionizing radiation, high intrinsic contrast, numerous techniques within one method and 3D reconstruction.

KEYWORDS: magnetic resonance imaging, perfusion, tissue characterization.

CITATION: *Cardiol Croat.* 2014;9(3-4):93.

Received: 15th Feb 2014

*Address for correspondence: Thalassoterapija Opatija, Maršala Tita 188/1, HR-51410 Opatija, Croatia.

Phone/Fax: +385 51 202 724

E-mail: viktor.persic@ri.t-com.hr

Literature

1. Nagel E, Lehmkuhl HB, BockschW, et al. Noninvasive diagnosis of ischemia-induced wall motion abnormalities with the use of high-dose dobutamine stress MRI: comparison with dobutamine stress echocardiography. *Circulation.* 1999;99:763-70.
2. Lima JA, Desai MY. Cardiovascular magnetic resonance imaging: current and emerging applications. *J Am Coll Cardiol.* 2004; 44:1164-71.
3. Lockie T, Ishida M, Perera D, et al. High-resolution magnetic resonance myocardial perfusion imaging at 3.0-Tesla to detect hemodynamically significant coronary stenoses as determined by fractional flow reserve. *J Am Coll Cardiol.* 2011;57:70-5.
4. Schwitler J, Wacker CM, Wilke N, et al. MR-IMPACTII: Magnetic Resonance Imaging for Myocardial Perfusion Assessment in Coronary artery disease Trial: perfusion-cardiac magnetic resonance vs. single-photon emission computed tomography for the detection of coronary artery disease: a comparative multicentre, multivendor trial. *Eur Heart J.* 2012; 34:775-81.
5. Kwong RY, Chan AK, Brown KA, et al. Impact of unrecognized myocardial scar detected by cardiac magnetic resonance imaging on event-free survival in patients presenting with signs or symptoms of coronary artery disease. *Circulation* 2006; 113:2733-43.