

**Randomized comparison of
intraaortic balloon counterpulsation
versus
optimal medical therapy in addition to early
revascularization in acute myocardial infarction
complicated by cardiogenic shock**

Holger Thiele, MD

Uwe Zeymer, MD; Franz-Josef Neumann, MD; Miroslaw Ferenc,
MD; Hans-Georg Olbrich, MD; Jörg Hausleiter, MD; Gert Richardt, MD;
Marcus Hennersdorf, MD; Klaus Empen, MD; Georg Fuernau, MD; Steffen Desch, MD;
Ingo Eitel, MD; Rainer Hambrecht, MD; Jörg Fuhrmann, MD; Michael Böhm, MD;
Henning Ebelt, MD; Steffen Schneider, PhD;
Gerhard Schuler, MD; Karl Werdan, MD

on behalf of the **IABP-SHOCK II Trial** Investigators

University of Leipzig – Heart Center

Disclosures

Funding:

German Research Foundation

German Heart Research Foundation

German Cardiac Society

Arbeitsgemeinschaft Leitende Kardiologische Krankenhausärzte

University of Leipzig – Heart Center

Unrestricted grant by:

Maquet Cardiopulmonary AG, Hirrlingen, Germany

Teleflex Medical, Everett, MA, USA

Potential Conflict of Interest:

Research Funding:

Terumo, Lilly, Maquet Cardiovascular, Teleflex Medical

Consulting:

Maquet Cardiovascular, Lilly

Speaker Honoraria:

Lilly, Astra Zeneca, Daiichi Sankyo, Boehringer Ingelheim, Maquet Cardiovascular, Medicines Company

History:

1962 Animal studies
Moulopoulos et al. Am Heart J 1962;63:669-675

1968 First clinical description in shock
Kantrowitz et al. JAMA 1968;203:135-140

1973 Hemodynamic effects in shock,
Mortality unchanged
Scheidt et al. NEJM 1973;288:979-984

> 40 years > 1 Million patients treated, low complication rate,
Benchmark registry
Ferguson et al. JACC 2001;38:1456-1462



IABP in AMI complicated by cardiogenic shock

ESC



Class IC


ACC/AHA




Class IB

Van de Werf et al. Eur Heart J 2008;29:2909-2945
Wijns et al. Eur Heart J 2010;31:2501-2555
Antman et al. Circulation 2004;110:82-292

Mortality IABP vs no IABP - Metaanalysis



IABP-Use in Cardiogenic Shock



Study Sites and Organisation



DSMB:

Kurt Huber
Ferenc Follath
Bernhard Maisch
Johannes Haerting


Steering committee:

Holger Thiele
Karl Werdan
Uwe Zeymer
Gerhard Schuler

Support + Patronage:



IABP-Shock-II Trial – Timelines



Sample Size

- Estimated 12% absolute difference in survival rates
- Sequential statistical design with 2 interim analyses (33% and 66% of patients)
- Significance level 0.0005 at 1st or 0.014 at 2nd interim analysis.
Final analysis at α -level 0.044 → 564 patients
- To compensate losses in follow-up and putative center effect → 600 patients

Primary Study Endpoint:

30-day all-cause mortality


Secondary Study Endpoints:

- Hemodynamic parameters (mean BP, heart rate pre and post revascularization)
- Serum-lactate (every 8 h for 48 h)
- SAPS-2 Score
- Serial creatinine-level and creatinine-clearance (Cockcroft-Gault-formula)
- Inflammatory reaction (CRP)

Process of care

- Time until hemodynamic stabilization
- Catecholamine dose and duration
- Requirement for LVAD-implantation or HTx
- Requirement for renal replacement therapy
- Length of ICU-stay
- Length of mechanical ventilation
- Mortality after 6 and 12 months

Trial Flow and Treatment




	IABP (n=301)	Control (n=299)
Age (years); median (IQR)	70 (58-78)	69 (58-76)
Male sex; n (%)	202 (67.1)	211 (70.6)
Current Smoking; n/total (%)	96/295 (32.5)	108/299 (36.1)
Hypertension; n/total (%)	213/296 (72.0)	199/299 (66.6)
Hypercholesterolemia; n/total (%)	122/295 (41.4)	105/299 (35.1)
Diabetes mellitus; n/total (%)	105/297 (35.4)	90/299 (30.1)
Prior myocardial infarction; n/total n (%)	71/300 (23.7)	61/299 (20.4)
Fibrinolysis < 24 h before randomization; n/total (%)	28/301 (9.3)	20/299 (6.7)
STEMI/LBBB; n/total (%)	200/300 (66.7)	212/298 (71.1)
NSTEMI; n/total (%)	96/300 (32.0)	81/298 (27.2)
Resuscitation before randomization; n/total (%)	127/301 (42.2%)	143/299 (47.8)
Signs of impaired organ perfusion; n/total (%)		
Altered mental status	215/300 (71.7)	232/299 (77.6)
Cold, clammy skin and extremities	257/300 (85.7)	245/299 (81.9)
Oliguria	90/300 (30.0)	99/299 (33.1)
Serum lactate >2.0 mmol/l	226/300 (75.3)	218/298 (73.2)
Creatinine clearance (ml/min); median (IQR)	60.7 (43.4-86.6)	56.8 (39.7-78.1)
Infarct related artery; n/total (%)		
LAD	132/293 (45.1)	121/293 (41.3)
LCX	55/293 (18.8)	57/293 (19.5)
RCA	73/293 (24.9)	79/293 (27.0)
Left main	26/293 (8.9)	28/293 (9.6)
Bypass graft	7/293 (2.4)	8/293 (2.7)
Multivessel disease; n/total (%)	235/296 (79.4)	228/293 (77.9)
Left ventricular ejection fraction (%); median (IQR)	35 (25-45)	35 (25-45)

Variable	IABP (n=301)	Control (n=299)	p
Primary PCI; n/total (%)	287/301 (95.3)	288/299 (96.3)	0.55
Stent implanted; n/total (%)	273/301 (90.7)	266/299 (89.0)	0.48
Drug-eluting stent; n/total (%)	126/301 (41.9)	123/299 (41.1)	0.86
Immediate PCI of non-culprit lesions; n/total (%)	90/301 (29.9)	81/299 (27.1)	0.45
Immediate bypass surgery; n/total (%)	8/301 (2.7)	10/299 (3.3)	0.62
Staged bypass surgery; n/total (%)	3/301 (1.0)	4/299 (1.3)	0.72
Active left ventricular assist device; n/total (%)	11/301 (3.7)	22/299 (7.4)	0.053
Mild hypothermia; n/total (%)	106/301 (35.2)	120/299 (40.1)	0.21
Mechanical ventilation; n/total (%)	240/301 (79.7)	252/299 (84.3)	0.15
Mechanical ventilation duration (days); median (IQR)	3.0 (1.0-8.0)	3.0 (1.0-8.0)	0.44
ICU treatment (days); median (IQR)	6.0 (3.0-12.0)	6.0 (3.0-13.0)	0.34
Renal replacement therapy; n/total (%)	62/301 (20.6)	47/299 (15.7)	0.12
Catecholamines ($\mu\text{g}/\text{kg}$ per minute); median (IQR)			
Dopamine	4.1 (2.9-7.7)	4.2 (3.6-8.3)	0.76
Norepinephrine	0.3 (0.1-1.2)	0.4 (0.1-1.1)	0.73
Epinephrine	0.3 (0.1-1.3)	0.3 (0.2-1.4)	0.59
Dobutamine	10.2 (4.9-20.6)	9.0 (4.8-17.6)	0.25
Duration of catecholamines (days), median (IQR)	3.0 (1.0-5.0)	3.0 (1.0-6.0)	0.81
Time - hemodynamic stabilization (days); median (IQR)	3.0 (1.0-5.0)	3.0 (1.0-6.0)	0.50


Simplified Acute Physiology Score-II



Renal Function (eGFR)




Serum Lactate




Inflammatory Reaction (CRP)



Primary Study Endpoint (30-Day Mortality)




Subgroups (30-Day Mortality)



	IABP (n=300)	Control (n=298)	P
Stroke in-hospital n/total (%)	2/300 (0.7)	5/298 (1.7)	0.28
GUSTO bleeding; n/total n (%)			
Life-threatening/severe	10/300 (3.3)	13/298 (4.4)	0.51
Moderate	52/300 (17.3)	49/298 (16.4)	0.77
Peripheral ischemic complication requiring intervention; n/total n (%)	13/300 (4.3)	10/298 (3.4)	0.53
Sepsis; n/total n (%)	47/300 (15.7)	61/298 (20.5)	0.15


Randomized Studies in Cardiogenic Shock



Summary + Conclusions

- IABP support in cardiogenic shock is safe without significant inherent complications.
- However, IABP support did not reduce 30-day mortality in this large, randomized, multicenter trial in cardiogenic shock patients complicating myocardial infarction undergoing early revascularization.
- The primary study endpoint results are supported by a lack of benefit in secondary endpoints.

Acknowledgement and Thank You



IABP-SHOCK II Investigators from 37 German Sites

Steering Committee

H. Thiele (Chair)
G. Schuler
K. Werdan
U. Zeymer

Funding

DFG
DSHF
DGK
ALKK
University of Leipzig –
Heart Center
Maquet Cardiovascular
Teleflex Medical

DSMB

J. Haerting (Chair)
F. Follath
K. Huber
B. Maisch

Study Sites

Leipzig: H. Thiele, G. Schuler, S. Desch, G. Fuernau, I. Eitel, S. de Waha, S. Wetzel, T. Pausewang, A. Leuschner
Bad Krozingen: F.-J. Neumann, M. Ferenc
Langen: H.-G. Olbrich, H.-B. Hopf
Munich German Heart Center: J. Hausleiter, A. de Waha, M. Orban, C. Lennerz, M. Seyfarth
Bad Segeberg: G. Richardt, B. Schwarz, M. Abdel-Wahab, R. Toelg, V. Geist, M. Bahnsen-Maaß
Heilbronn: M. Hennersdorf, U. Rieman, J. Graf, A. Kuhn, D. Scharpf
Greifswald: K. Empen, S. Felix
Halle (Saale): K. Werdan, H. Ebelt, A. Schlitt, M. Buerke
Bremen: R. Hambrecht, E. Fiehn, A. Fach
Ludwigshafen: U. Zeymer, A. K. Gitt, B. Mark, R. Winkler
Bad Berka: B. Lauer, J. Fuhrmann
Homburg/Saar: M. Böhm, A. Link
Jena: H.R. Figulla, M. Ferrari, C. Jung, S. Utschig
Lübeck: V. Kurowski, S. Wolfrum, P.W. Radke, H. Schunkert
Hennigsdorf: H.-H. Minden
Magdeburg: R. C. Braun-Dullaeus, F. Walz, A. Schmeißer
Cottbus – Heart Center: J. Krülls-Münch, K. Rochor
Dresden: R. H. Strasser, G. Simonis
Würzburg: S. Maier, G. Ertl
Munich – Neuperlach: H. Mudra, M. Hug
Regensburg: P. Bomba, P. Sick
Berlin – Benjamin Franklin: C. Tschöpe, H.-P. Schutlheiss
Berlin – Vivantes am Urban: H. Roth, D. Andresen
Cottbus – Carl-Thiem-Klinikum: C. Kurek, J. Krülls-Münch
Rostock: C. Nienaber, H. Ince, H. Schneider
Bad Nauheim: C. Hamm, H. Möllmann
Erfurt: H. Lapp
Berlin – Charité Mitte: G. Baumann, F. Knebel
Munich – Großhadern: W. Franz
Berlin – Vivantes AVK: H. Schühlen, L. U. Sttracke
Weiden: R. H. Schwinger, H. Bäuml
Fulda: V. Schächinger
Essen: M. Lichtenberg
Unna: D. Gießmann
Merseburg: R. Prondzinsky
Detmold: U. Tebbe
Villingen-Schwenningen: R. Birkemeyer

CEC

H. Thiele
U. Zeymer

CRO IHF Ludwigshafen

S. Schneider (Statistics chair)
T. Oarrak (Statistics)
U. Zeymer
K. Vonderschmitt
Z. Alkisoglu
S. Frey
B. Messemer

The NEW ENGLAND JOURNAL of MEDICINE

ORIGINAL ARTICLE

Intraaortic Balloon Support for Myocardial Infarction with Cardiogenic Shock

Holger Thiele, M.D., Uwe Zeymer, M.D., Franz-Josef Neumann, M.D., Miroslaw Ferenc, M.D., Hans-Georg Olbrich, M.D., Jörg Hausleiter, M.D., Gert Richardt, M.D., Marcus Hennersdorf, M.D., Klaus Empen, M.D., Georg Fuernau, M.D., Steffen Desch, M.D., Ingo Eitel, M.D., Rainer Hambrecht, M.D., Jörg Fuhrmann, M.D., Michael Böhm, M.D., Henning Ebel, M.D., Steffen Schneider, Ph.D., Gerhard Schuler, M.D., and Karl Werdan, M.D., for the IABP-SHOCK II Trial Investigators*