

Multimodality imaging-based diagnosis of cardiac transthyretin amyloidosis

Koraljka Benko¹,
Tamara Hlača Caput¹,
Saša Matulić¹,
Danijel Premuš²,
Ivana Grgić Romić¹,
Ana Hrelja¹,
Iva Žuža¹,
Teodora Zaninović
Jurjević^{1*}

¹University Hospital Centre
Rijeka, Rijeka, Croatia

²Hospital for Medical
Rehabilitation of Heart
and Lung Diseases
and Rheumatism
"Thalassotherapia-Opatija",
Opatija, Croatia

KEYWORDS: amyloidosis, transthyretin, heart failure.

CITATION: Cardiol Croat. 2023;18(5-6):170. | <https://doi.org/10.15836/ccar2023.170>

***ADDRESS FOR CORRESPONDENCE:** Teodora Zaninović Jurjević, Klinički bolnički centar Rijeka, Tome Strišića 3, HR-51000 Rijeka, Croatia. / Phone: +385-51-407225 / E-mail: teodorazj@gmail.com

ORCID: Koraljka Benko, <https://orcid.org/0000-0001-7556-0860> • Tamara Hlača Caput, <https://orcid.org/0000-0002-7749-0031>
Saša Matulić, <https://orcid.org/0009-0004-0730-1380> • Danijel Premuš, <https://orcid.org/0000-0002-6806-2027>
Ivana Grgić Romić, <https://orcid.org/0000-0002-0035-4445> • Ana Hrelja, <https://orcid.org/0009-0009-0892-5220>
Iva Žuža, <https://orcid.org/0000-0002-5542-0091> • Teodora Zaninović Jurjević, <https://orcid.org/0000-0001-8359-3910>

Introduction: Transthyretin amyloidosis cardiomyopathy (ATTR-CM) is a rare, progressive, life-threatening, hereditary disorder induced by a misfolded precursor protein, caused by mutations in the transthyretin gene. ATTR-CM is a challenging disease to recognize in early stages owing to its multisystem and nonspecific manifestations¹⁻³.

Case report: 65-year-old patient was hospitalized for the second time to our Clinic for Cardiovascular diseases in April 2022. Previous hospitalization happened 5 years ago and there was no ambulatory cardiology check-up between hospitalizations. He had progressive dyspnea on mild exertion and lower extremity edema. Therapy at home was furosemide occasionally. In past medical history he had left carpal tunnel surgery and Covid 19 infection in April 2022. On arrival blood pressure was 140/80 mmHg, pulse 100 beats per minute, absolutely arrhythmic. Basal weakened breath sounds, and leg edema were present. Troponin and NT pro BNP were elevated (91 ng/l and 3222 ng/l). 12-lead electrocardiogram showed peripheral micro-voltage, atrial fibrillation with ventricular rate around 110 bpm. Biatrial enlargement, increased left and right wall thickness, thickened papillary muscles, mildly reduced left ventricular ejection fraction and mono-phasic transmittal flow was found on transthoracic echocardiography. Testing for Fabry disease was negative. Cardiac magnetic resonance (CMR) found morphological changes and the pattern of contrast accumulation that suggested cardiac amyloidosis. Immunofixation electrophoresis showed no monoclonal (M) spike, gammopathy was unlikely. A biopsy of the buccal mucosa was performed, no amyloid deposits were found. Bone scintigraphy found accumulation of labelled hydroxydiphosphonate (HDP) that was visible in the myocardium, which points to ATTR-CM. Genetic testing is in progress.

Conclusion: ATTR-CM requires a high index of suspicion, and it should be suspected in patients with LV hypertrophy and heart failure. The diagnosis of cardiac amyloidosis requires a combination of multi-modality imaging including echocardiography, CMR and scintigraphy. An imaging modality that can accurately diagnose ATTR-CM without the need for invasive cardiac biopsy is nuclear scintigraphy using bone-avid radio-tracers⁴. Timely diagnosis is important since the treatment is possible and improves prognosis in these patients.

RECEIVED:
March 26 2023

ACCEPTED:
March 29, 2023



LITERATURE

1. Planté-Bordeneuve V, Said G. Familial amyloid polyneuropathy. *Lancet Neurol*. 2011 Dec;10(12):1086-97. [https://doi.org/10.1016/S1474-4422\(11\)70246-0](https://doi.org/10.1016/S1474-4422(11)70246-0)
2. Olsson M, Hellman U, Planté-Bordeneuve V, Jonasson J, Lång K, Suhr OB. Mitochondrial haplogroup is associated with the phenotype of familial amyloidosis with polyneuropathy in Swedish and French patients. *Clin Genet*. 2009 Feb;75(2):163-8. <https://doi.org/10.1111/j.1399-0004.2008.01097.x>
3. Gillmore JD, Dmy T, Fontana M, Hutchinson M, Lachmann HJ, Martinez-Naharro A, et al. A new staging system for cardiac transthyretin amyloidosis. *Eur Heart J*. 2018 Aug 7;39(30):2799-2806. <https://doi.org/10.1093/eurheartj/ehx589>
4. Perugini E, Guidalotti PL, Salvi F, Cooke RM, Pettinato C, Riva L, et al. Noninvasive etiologic diagnosis of cardiac amyloidosis using 99mTc-3,3'-diphosphono-1,2-propanodicarboxylic acid scintigraphy. *J Am Coll Cardiol*. 2005 Sep 20;46(6):1076-84. <https://doi.org/10.1016/j.jacc.2005.05.073>