

Mitral annular disjunction and cardiac magnetic resonance

 **Vesna Pehar Pejčinović¹,**
 **Iva Uravić Bursać^{1,*},**
 **Dijana Travica Samsa¹,**
 **Marijana Rakić¹,**
 **Damir Rajčević¹,**
 **Viktor Peršić^{1,2}**

¹Hospital for Medical Rehabilitation of Heart and Lung Diseases and Rheumatism "Thalassotherapia-Opatija", Opatija, Croatia

²University of Rijeka, Faculty of Medicine, Rijeka, Croatia

KEYWORDS: cardiac magnetic resonance, mitral annular disjunction, prolapse.

CITATION: *Cardiol Croat.* 2023;18(5-6):173. | <https://doi.org/10.15836/ccar2023.173>

***ADDRESS FOR CORRESPONDENCE:** Iva Uravić Bursać, "Thalassotherapia-Opatija", M. Tita 180, HR-51410 Opatija, Croatia. / Phone: + 385-51-202-600 / E-mail: bursaciva@gmail.com

ORCID: Vesna Pehar Pejčinović, <https://orcid.org/0000-0002-8921-7999> • Iva Uravić Bursać, <https://orcid.org/0000-0002-1050-0135> • Dijana Travica Samsa, <https://orcid.org/0000-0001-6238-3738> • Marijana Rakić, <https://orcid.org/0000-0002-7462-054X> • Damir Rajčević, <https://orcid.org/0000-0001-9743-9201> • Viktor Peršić, <https://orcid.org/0000-0003-4473-5431>

Mitral annular disjunction (MAD) is the improper systolic acceptance of the posterior leaflet to the atrial wall¹. First line to the diagnosis is transthoracic echocardiography in the long axis view on the mitral valve using highest frame rate or sagittal view on cardiac magnetic resonance (CMR)¹. The distance between mitral-annulus and systolic bulge of ventricular myocardium range 5-10 mm is diagnostic and it is associated with morphological and functional remodeling of the left ventricular myocardium^{1,2}. Prevalence of MAD varies due to the different cut-offs, imaging modalities and ec, and in a general population, is 8.7%³. In patients with mitral valve prolapse the prevalence of MAD is 20-58%¹. While MAD with prolapse is common and associated with ventricular arrhythmias, isolated MAD without prolapse, has been described on CMR^{2,3}. CMR can also identify prolapse using cine images, but the most important advantage of this technique is risk stratification of arrhythmias due to incremental prognostic value of late gadolinium enhancement over mitral valve prolapse severity². Several studies suggested association between late gadolinium enhancement (LGE) at the mid wall of papillary muscles and inferobasal region of left ventricle with complex arrhythmias¹⁻⁴. The origin of ventricular arrhythmia can be deduced according to the distribution of LGE on CMR^{1,2}. CMR should be done in all patients who survived sudden cardiac death or sustained ventricular arrhythmia before devices, to clarify the etiology and also in patients with unexplained syncope or nonsustained ventricular arrhythmia, to assessment of left ventricle size and function, severity of mitral regurgitation, leaflet thickness and also when is poor echocardiographic window¹. CMR imaging provides excellent morphological information and helps in the assessment of fibrosis⁴. Myocardial fibrosis determined according to LGE at CMR was associated with adverse outcome in patients with mitral valve prolapse without moderate-to-severe mitral regurgitation or left ventricular dysfunction².

RECEIVED:
March 4 2023

ACCEPTED:
March 29, 2023



LITERATURE

1. Sabbag A, Essayagh B, Barrera JDR, Basso C, Berni A, Cosyns B, et al. EHRA expert consensus statement on arrhythmic mitral valve prolapse and mitral annular disjunction complex in collaboration with the ESC Council on valvular heart disease and the European Association of Cardiovascular Imaging endorsed by the Heart Rhythm Society, by the Asia Pacific Heart Rhythm Society, and by the Latin American Heart Rhythm Society. *Europace*. 2022 Dec 9;24(12):1981-2003. <https://doi.org/10.1093/europace/euac125>
2. Figliozzi S, Georgiopoulos G, Lopes PM, Bauer KB, Moura-Ferreira S, Tondi L, et al. Myocardial Fibrosis at Cardiac MRI Helps Predict Adverse Clinical Outcome in Patients with Mitral Valve Prolapse. *Radiology*. 2023 Jan;306(1):112-121. <https://doi.org/10.1148/radiol.220454>
3. Bennett S, Tafuro J, Duckett S, Appaji A, Khan JN, Heatlie G, et al. Definition, prevalence, and clinical significance of mitral annular disjunction in different patient cohorts: A systematic review. *Echocardiography*. 2022 Mar;39(3):514-523. <https://doi.org/10.1111/echo.15299>
4. Kulkarni AA, Chudgar PD, Burkule NJ, Kamat NV. Mitral Annulus Disjunction and Arrhythmic Mitral Valve Prolapse: Emerging Role of Cardiac Magnetic Resonance Imaging in the Workup. *Indian J Radiol Imaging*. 2022 Aug 30;32(4):576-581. <https://doi.org/10.1055/s-0042-1754357>