

Pulmonary vein isolation by cryoablation at the General Hospital Slavonski Brod

Alenka Tuličić-Mihelčić*,
Renata Valenčak,
Saša Presežnik,
Marina Stanković

General Hospital "Dr Josip Benčević", Slavonski Brod, Croatia

KEYWORDS: atrial fibrillation, pulmonary vein isolation, cryoablation.

CITATION: Cardiol Croat. 2026;21(1-2):31. | <https://doi.org/10.15836/ccar2026.31>

***ADDRESS FOR CORRESPONDENCE:** Alenka Tuličić-Mihelčić, Opća bolnica "Dr Josip Benčević", Andrije Štampara 42, HR-35000 Slavonski Brod, Croatia. / Phone: +385-98-771-551 / E-mail: alenkaveyron@gmail.com

ORCID: Alenka Tuličić-Mihelčić, <https://orcid.org/0000-0002-4745-1066> • Renata Valenčak, <https://orcid.org/0000-0001-6323-462X> Saša Presežnik, <https://orcid.org/0009-0006-8893-6365> • Marina Stanković, <https://orcid.org/0000-0002-8717-0357>

Atrial fibrillation (AF) is the most common cardiac arrhythmia in the general population. The global prevalence of AF is estimated to be between 2% and 4%, with a continued increase in the number of patients expected¹. AF is associated with increased overall mortality and morbidity, as well as a higher risk of stroke and heart failure². The gold standard in the treatment of AF is pulmonary vein isolation. At the General Hospital Slavonski Brod, the first pulmonary vein isolation using cryoablation was performed on February 13, 2019. Since then, a total of 186 procedures have been conducted. The procedure is performed by an interventional cardiologist/electrophysiologist. At the beginning of the procedure, local anesthetic is applied to both groin regions. The left femoral vein is punctured, and an introducer sheath is inserted through which a line for measuring activated clotting time, an invasive pressure monitoring system, and intracardiac echocardiography (ICE) are placed. Through the puncture of the right femoral artery, an introducer with an inserted needle for transseptal puncture is positioned, enabling access to the left atrium and pulmonary vein ostia. A cryoballoon is then inserted through the same introducer, and ablation of the pulmonary vein ostia is performed using liquid nitrogen. ICE is removed, and a decapolar electrophysiological catheter is placed to monitor the function of the phrenic nerve via electrical stimulation. If, at the end of the procedure, the patient is still in atrial fibrillation, synchronized cardioversion is performed¹.

As with all invasive cardiology procedures, aseptic techniques, sterility protocols, hemodynamic monitoring, analgesia and sedation, coagulation time measurement, and fluoroscopy are all employed, alongside other standard procedural steps. The medical team includes two nurses and a radiology technologist. Potential complications include allergic reactions to the contrast agent, puncture site complications such as bleeding, hematoma, or infection, phrenic nerve injury, esophageal injury, cardiac perforation, pulmonary vein damage, or stroke. This is why the procedure requires thorough preparation and a well-trained, experienced team².

RECEIVED:
October 15, 2025

ACCEPTED:
November 14, 2025



LITERATURE

- Hindricks G, Potpara T, Dagres N, Arbelo E, Bax JJ, Blomström-Lundqvist C, et al; ESC Scientific Document Group. 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS): The Task Force for the diagnosis and management of atrial fibrillation of the European Society of Cardiology (ESC) Developed with the special contribution of the European Heart Rhythm Association (EHRA) of the ESC. Eur Heart J. 2021 Feb 1;42(5):373-498. <https://doi.org/10.1093/eurheartj/ehaa612>
- January CT, Wann LS, Alpert JS, Calkins H, Cigarroa JE, Cleveland JC Jr, et al; ACC/AHA Task Force Members. 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines and the Heart Rhythm Society. Circulation. 2014 Dec 2;130(23):e199-267. <https://doi.org/10.1161/CIR.0000000000000041>